

HEALTH TECHNOLOGY BRIEFING OCTOBER 2020

Lisocabtagene maraleucel for transplant-eligible and ineligible, relapsed or refractory, aggressive B-cell non-Hodgkin lymphoma - second-line

NIHRIO ID	24260	NICE ID	10383
Developer/Company	Celgene Ltd	UKPS ID	655777

Licensing and market availability plans	Currently in phase III clinical trials.
--	---

SUMMARY

Lisocabtagene maraleucel is in clinical development for the treatment of adult patients with high-risk, relapsed or refractory aggressive B-cell non-Hodgkin lymphoma (NHL). In NHL, the affected lymphocytes start to multiply abnormally and begin to collect in certain parts of the body such as the lymph nodes. Aggressive or high-grade lymphomas tend to grow very quickly. Relapsed means the lymphoma has responded to treatment but then returns, and refractory means the lymphoma has not responded to the initial treatment. Treatment options for relapsed/refractory aggressive B-cell lymphoma are limited.

Lisocabtagene maraleucel is administered intravenously. It contains the patient's white blood cells (T-cells) that have been modified genetically in the laboratory so that they make a protein called chimeric antigen receptor (CAR). The CAR T-cells attach to a particular protein called CD19 on the surface of cancer cells resulting in the death of these cells. If licensed, lisocabtagene maraleucel will offer a treatment option for adult patients with high-risk, relapsed or refractory aggressive B-cell NHL.

PROPOSED INDICATION

Second-line treatment for adult patients with high-risk, relapsed or refractory aggressive B-cell non-Hodgkin lymphoma (NHL).^{1,2}

TECHNOLOGY

DESCRIPTION

Lisocabtagene maraleucel (Liso-cel) is a chimeric antigen receptor (CAR) T-cell therapy designed to target CD19 which is a surface glycoprotein expressed during normal B-cell development and maintained following malignant transformation of B cells. Lisocabtagene maraleucel aims to target CD-19 expressing cells through a CAR construct that includes an anti-CD19 single chain variable fragment targeting domain for antigen specificity, a transmembrane domain, a 4-1BB costimulatory domain hypothesized to increase T-cell proliferation and persistence, and a CD3-zeta T-cell activation domain.³ Studies suggest CD-19 CAR-T cell-based immunotherapies effectively recognize and kill CD-19-positive target cells and result in a high clinical response rate in the treatment of refractory B-cell malignancies.⁴

Lisocabtagene maraleucel is in clinical development for the treatment of adult patients with high-risk, relapsed or refractory aggressive B-cell NHL. In the phase III clinical trial (TRANSFORM; NCT03575351), patients will receive lymphodepleting chemotherapy with intravenous (IV) fludarabine (30 mg/m²/day for 3 days) plus cyclophosphamide IV (300 mg/m²/day for 3 days) concurrently followed by lisocabtagene maraleucel infusion.¹ In the phase II clinical trial (TRANSCEND-PILOT-017006; NCT03483103) participants will receive a single dose of lisocabtagene maraleucel (100x10⁶ CAR+ T cells) by IV injection on day 1 (between 2 and 7 days following the completion of lymphodepleting chemotherapy).²

INNOVATION AND/OR ADVANTAGES

An advantage of CAR T-cell therapy is the short treatment time needed because the therapy is administered with a single infusion. The recovery time is also much more rapid compared to stem cell transplants where aggressive chemotherapy is also used.⁵

Lisocabtagene maraleucel has a highly controlled manufacturing process that, unlike other CAR T cell therapies, enables administration of a defined composition with a precise dose of CD8+ and CD4+ CAR T cells. The manufacturing process and control strategy reduces the risk of complete manufacturing failure and minimises product variability in quality attributes, which may contribute to an improved safety and efficacy profile.⁶ Lisocabtagene maraleucel appears to exhibit a safer toxicity profile compared to axicabtagene ciloleucel and tisagenlecleucel, which are the other CD19-targeted CAR-T cell therapies approved in the treatment of relapsed/refractory, aggressive non- NHL, and has the potential for outpatient administration.⁷

DEVELOPMENT STATUS AND/OR REGULATORY DESIGNATIONS

Lisocabtagene maraleucel does not currently have Marketing Authorisation in the EU/UK for any indication. However, a Marketing Authorisation Application was submitted in June 2020 for adults with relapsed/refractory DLBCL, PMBCL and FL3B after at least two prior therapies.

Lisocabtagene maraleucel was granted an orphan designation for the treatment of primary mediastinal large B-cell lymphoma by the EMA in November 2018.⁸

Lisocabtagene maraleucel is currently in phase II clinical development for the treatment of other forms of lymphoma and lymphocytic leukaemia.⁹

PATIENT GROUP

DISEASE BACKGROUND

Lymphoma is a cancer of the lymphatic system. The lymphatic system is a system of lymphatic vessels and lymph nodes that run throughout the body. Tissue fluid called lymph circulates around the body in these vessels and flows through the lymph nodes. The lymphatic system is an important part of our immune system that plays a role in fighting bacteria and other infections and it tries to destroy old or abnormal cells, such as cancer cells. In people who have lymphoma, white blood cells divide abnormally before they are fully mature. This means they cannot fight infection as normal white blood cells do and they begin to collect in the lymph nodes or other places such as the bone marrow or spleen. They can then grow into tumours and begin to cause problems in the lymphatic system or in the organ in which they are growing.¹⁰

There are two main types of lymphoma: Hodgkin lymphoma and NHL. In NHL, the affected lymphocytes start to multiply abnormally and begin to collect in certain parts of the lymphatic system, such as the lymph nodes.¹¹ NHL is divided into more than 30 types, classified based on the type of lymphocyte involved: B lymphocytes (B cells) or T lymphocytes (T cells). NHL can be further classified by other factors such as whether it is aggressive (fast-growing) or indolent (slow-growing).¹² Relapsed lymphoma is when the lymphoma comes back after successful treatment and a period of remission (no evidence of lymphoma on tests and scans). Refractory lymphoma is when the lymphoma does not respond well to the first choice of treatment.¹³

The most common symptom of NHL is a painless swelling in a lymph node, usually in the neck, armpit or groin. Other more general symptoms can night sweats, unintentional weight loss, a high temperature, breathlessness or persistent itching of the skin all over the body.¹⁴ NHL is caused by a mutation in the DNA of lymphocytes although the exact reason for this mutation isn't known. Risk factors for developing NHL include having: a medical condition that weakens the immune system, medical treatment that weakens the immune system, autoimmune conditions, been infected with the Epstein-Barr virus, or Helicobacter pylori, received chemotherapy or radiotherapy for an earlier cancer and coeliac disease.¹⁵

CLINICAL NEED AND BURDEN OF DISEASE

According to the Haematological Malignancy Research Network, in the UK in 2016, there was estimated to be 5,510 people with diffuse large B-cell lymphoma (DLBCL) and 2,220 with follicular lymphoma.¹⁶

Overall, for NHL, European age-standardised incidence rates are projected to decrease by 2.75% from 32.45 per 100,000 in 2014 to 31.56 per 100,000 in 2035 in males, and by 3.32% from 22.67 per 100,000 in 2014 to 21.92 per 100,000 in 2035 in females.¹⁷

In England, in 2017, there were a total of 4,348 registrations of deaths due to NHL (ICD-10 code: C82-C85).¹⁸ The age-standardised one-year and five-year survival rates for NHL (all subtypes combined) in England over the period 2013 and 2017, followed up to 2018, show that 78.4% of men are expected to survive for at least 1 year, and 63.7% are expected to survive for 5 years. The survival rates for women are slightly higher with 80.6% expected to survive for at least 1 year and 68.1% expected to survive for 5 years.¹⁹

PATIENT TREATMENT PATHWAY

TREATMENT PATHWAY

Treatment of NHL is determined by a multidisciplinary team which usually includes a haematologist, cancer nurse, a pathologist, a radiotherapy specialist, a pharmacist and radiologist. Treatment depends on the type and grade of NHL, the stage of NHL, general health of the patient and the patient's age.²⁰

The main types of treatment for NHL are chemotherapy, targeted cancer drugs and radiotherapy. Other treatment types include stem cell transplant and surgery.²⁰

CURRENT TREATMENT OPTIONS

Treatment depends on the type and grade of NHL, the stage of NHL, general health of the patient and the patient's age.²⁰

The main types of treatment for NHL are chemotherapy, targeted cancer drugs and radiotherapy. Other treatment types include stem cell transplant and surgery. Some patients only have one type of treatment. Other patients may need a combination of treatments²⁰

PLACE OF TECHNOLOGY

If licensed, lisocabtagene maraleucel will provide a treatment option for adult patients with high-risk, relapsed or refractory aggressive B-cell NHL.

CLINICAL TRIAL INFORMATION

Trial	TRANSFORM , NCT03575351 , EudraCT 2018-000929-32 ; A Global Randomized Multicenter Phase 3 Trial of JCAR017 Compared to Standard of Care in Adult Subjects With High-risk, Second-line, Transplant-eligible Relapsed or Refractory Aggressive B-cell Non-Hodgkin Lymphomas (TRANSFORM) Trial phase III- Recruiting Location(s): EU countries (Incl UK): USA and Japan Primary completion date: January 2024
Trial design	Randomised, parallel assignment, open-label
Population	N=182 (planned); adults aged 18-75 years; diffuse large B-cell lymphoma; refractory or relapsed within 12 months from CD20 antibody and anthracycline containing first line therapy
Intervention(s)	Lisocabtagene maraleucel (IV)
Comparator(s)	Standard of care
Outcome(s)	Event-free survival (Time Frame: approximately 3 years) See trial record for full list of other outcomes.
Results (efficacy)	-
Results (safety)	-

Trial	TRANSCEND-PILOT, NCT03483103 ; A Phase 2 Study of Lisocabtagene (JCAR017) as Second-Line Therapy in Adult Patients With Aggressive B-cell NHL (017006) Phase II – Recruiting Location: United States Primary completion date: April 2021
Trial design	Single group assignment, open-label
Population	N=56 (planned); adults aged 18 years and older; confirmation of relapsed or refractory aggressive B-cell non-Hodgkin lymphoma; previous treatment with a single line of chemo-immunotherapy containing an anthracycline and a CD20 targeted agent
Intervention(s)	Lisocabtagene maraleucel (IV)
Comparator(s)	No comparator
Outcome(s)	Primary outcome measure: <ul style="list-style-type: none"> Overall response rate – anti-tumour activity (Time frame: through month 24) See trial record for full list of other outcomes.
Results (efficacy)	At data cut off, 25 patients had lymphodepleting chemotherapy followed by lisocabtagene maraleucel infusion. At median follow-up of 3.5 months the objective response rate was 80%. 48% of patients achieved complete response. ²¹
Results (safety)	72% of patients had grade ≥3 treatment emergent adverse events (TEAEs), 40% of which were cytopenias. No grade 5 adverse events (AEs) occurred within the first 30 days after lisocabtagene infusion. Five patients (20%) had cytokine release syndrome (CRS) and 3 (12%) had neurological events (NE). No grade 3 or 4 CRS was observed; 2 patients (8%) had grade 3 or 4 NEs. Five patients (20%) received tocilizumab and/or dexamethasone for CRS/NEs. ²¹

ESTIMATED COST

The estimated cost of Lisocabtagene maraleucel is not yet known.

RELEVANT GUIDANCE

NICE GUIDANCE

- NICE technology appraisal in development. Ibrutinib for treating relapsed or refractory follicular lymphoma (TA10223). Expected date of issue to be confirmed.
- NICE technology appraisal in development. Tafastiamab with lenalidomide for previously treated diffuse large B-cell lymphoma (TA10645). Expected date of issue to be confirmed.
- NICE technology appraisal in development. Axicabtagene ciloleucel for treating relapsed or refractory diffuse large B-cell lymphoma after 1 systemic therapy (TA10580). Expected date of issue to be confirmed.
- NICE technology appraisal in development. Polatuzumab vedotin with rituximab and bendamustine for treating relapsed or refractory diffuse large B-cell lymphoma (TA10463). Expected publication date: September 2020.

- NICE technology appraisal. Lenalidomide with rituximab for previously treated follicular lymphoma (TA627). April 2020.
- NICE technology appraisal. Rituximab for the treatment of relapsed or refractory stage III or IV follicular non-Hodgkin's lymphoma (TA137). February 2008.
- NICE guideline. Non-Hodgkin's lymphoma: diagnosis and management (NG52). 2016.
- NICE quality standard. Haematological cancers (QS150). 2017.

NHS ENGLAND (POLICY/COMMISSIONING) GUIDANCE

- No relevant guidance identified

OTHER GUIDANCE

- European Society of Medical Oncology (ESMO). Newly diagnosed and relapsed follicular lymphoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. 2016.²²
- European Society of Medical Oncology (ESMO). Diffuse large B-cell lymphoma (DLBCL): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. 2015.²³
- London Cancer. Guidelines for the management of non-Hodgkin's and Hodgkin's lymphoma in adults. 2014.²⁴

ADDITIONAL INFORMATION

REFERENCES

- 1 ClinicalTrials.gov. *A Global Randomized Multicenter Phase 3 Trial of JCAR017 Compared to Standard of Care in Adult Subjects With High-risk, Second-line, Transplant-eligible Relapsed or Refractory Aggressive B-cell Non-Hodgkin Lymphomas (TRANSFORM)*. Trial ID: NCT03575351. Status: Recruiting. Available from: <https://clinicaltrials.gov/ct2/show/NCT03575351> [Accessed 29 September 2020].
- 2 ClinicalTrials.gov. *A Phase 2 Study of Lisocabtagene Maraleucel (JCAR017) as Second-Line Therapy in Adult Patients With Aggressive B-cell NHL (017006)*. Trial ID: NCT03483103. Available from: <https://clinicaltrials.gov/ct2/show/NCT03483103> [Accessed 25 October 2020].
- 3 Bristol Myers Squibb. *Bristol Myers Squibb Provides Update on Biologics Licence Application (BLA) for Lisocabtagene Maraleucel (liso-cel)*. 2020. Available from: <https://news.bms.com/press-release/corporatefinancial-news/bristol-myers-squibb-provides-update-biologics-licence-applica> [Accessed 29 September 2020].
- 4 Informa Pharma Intelligence. *Lisocabtagene maraleucel*. 2019. Available from: <https://pharmastore.informa.com/product/lisocabtagene-maraleucel/> [Accessed 29 September 2020].
- 5 Rutgers Cancer Institute of New Jersey. *Advantages of CAR-T Cell Therapy*. 2020. Available from: <https://www.cinj.org/patient-care/advantages-car-t-cell-therapy> [Accessed 29 September 2020].
- 6 Ramsborg CG, Guptill P, Weber C, Christin B, Larson RP, Lewis K, et al. JCAR017 Is a Defined Composition CAR T Cell Product with Product and Process Controls That Deliver Precise Doses of CD4 and CD8 CAR T Cell to Patients with NHL. *Blood*. 2017;130(Supplement 1):4471-. Available from: https://doi.org/10.1182/blood.V130.Suppl_1.4471.4471.
- 7 Quintás-Cardama A. CD19 directed CAR T cell therapy in diffuse large B-cell lymphoma. *Oncotarget*. 2018;9(52):29843-4. Available from: <https://doi.org/10.18632/oncotarget.25688>.
- 8 European Medicines Agency. *Public summary of opinion on orphan designation: Lisocabtagene maraleucel for the treatment of primary mediastinal large B-cell lymphoma*. 2019. Available from: <https://www.ema.europa.eu/en/documents/orphan-designation/eu/3/18/2099-public-summary->

- [opinion-orphan-designation-lisocabtagene-maraleucl-treatment-primary_en.pdf](#) [Accessed 29 September 2020].
- 9 Clinicaltrials.gov. *Search for lisocabtagene studies: Phase II*. 2020. Available from: https://clinicaltrials.gov/ct2/results?intr=lisocabtagene&age_v=&gndr=&type=&rslt=&phase=1&Search=Apply [Accessed 29 September 2020].
- 10 Cancer Research UK. *What is non-Hodgkin Lymphoma?* 2018. Available from: <https://www.cancerresearchuk.org/about-cancer/non-hodgkin-lymphoma/about> [Accessed 29 September 2020].
- 11 National Health Service. *Overview: Non-Hodgkin lymphoma*. 2018. Available from: <https://www.nhs.uk/conditions/non-hodgkin-lymphoma/> [Accessed 29 September 2020].
- 12 Cancer Treatment Centers of America. *Non-Hodgkin lymphoma types*. 2020. Available from: <https://www.cancercenter.com/cancer-types/non-hodgkin-lymphoma/types> [Accessed 29 September 2020].
- 13 Lymphoma Action. *What happens if lymphoma relapses?* 2020. Available from: <https://lymphoma-action.org.uk/about-lymphoma-living-and-beyond-lymphoma/what-happens-if-lymphoma-relapses> [Accessed 29 September 2020].
- 14 National Health Service. *Non-Hodgkin lymphoma: Symptoms*. 2018. Available from: <https://www.nhs.uk/conditions/non-hodgkin-lymphoma/symptoms/> [Accessed 29 September 2020].
- 15 National Health Service. *Non-Hodgkin Lymphoma: Causes*. 2018. Available from: <https://www.nhs.uk/conditions/non-hodgkin-lymphoma/causes/> [Accessed 29 September 2020].
- 16 University of York. *HMRN Incidence*. 2020. Available from: <https://www.hmrn.org/statistics/incidence> [Accessed 29 September 2020].
- 17 Cancer Research UK. *Cancer incidence for common cancers*. 2020. Available from: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/incidence/common-cancers-compared#heading-Four> [Accessed 29 September 2020].
- 18 Office for National Statistics (ONS). *Death Registration Summary Statistics, England and Wales, 2017*. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/deathregistrationssummarytablesenglandandwalesreferencetables> [Downloaded 29 September 2020].
- 19 Office for National Statistics (ONS). *Cancer survival in England: adult, stage at diagnosis and childhood - patients followed up to 2018*. Available from: <https://www.ons.gov.uk/releases/cancersurvivalinenglandadultstageatdiagnosisandchildhoodpatientsfollowedupto2018> [Downloaded 29 September 2020].
- 20 Cancer Research UK. *Non-Hodgkin Lymphoma: Treatment decisions*. 2018. Available from: <https://www.cancerresearchuk.org/about-cancer/non-hodgkin-lymphoma/treatment/decisions> [Accessed 29 September 2020].
- 21 Sehgal AR, Hildebrandt G, Ghosh N, Godwin JE, Wagner-Johnston ND, Hoda D, et al. Lisocabtagene maraleucl (liso-cel) for treatment of second-line (2L) transplant noneligible (TNE) relapsed/refractory (R/R) aggressive large B-cell non-Hodgkin lymphoma (NHL): Updated results from the PILOT study. *Journal of Clinical Oncology*. 2020 2020/05/20;38(15_suppl):8040-. Available from: https://doi.org/10.1200/JCO.2020.38.15_suppl.8040.
- 22 Dreyling M, Ghielmini M, Rule S, Salles G, Vitolo U, Ladetto M. Newly diagnosed and relapsed follicular lymphoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2016 Sep;27(suppl 5):v83-v90. Available from: <https://doi.org/10.1093/annonc/mdw400>.
- 23 Tilly H, Gomes da Silva M, Vitolo U, Jack A, Meignan M, Lopez-Guillermo A, et al. Diffuse large B-cell lymphoma (DLBCL): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2015 Sep;26 Suppl 5:v116-25. Available from: <https://doi.org/10.1093/annonc/mdv304>.
- 24 London Cancer. *Guidelines for the management of Lymphoma*. 2014. Available from: <http://www.londoncancer.org/media/84337/london-cancer-lymphoma-radiotherapy-guidelines.pdf> [Accessed 29 September 2020].

NB: This briefing presents independent research funded by the National Institute for Health Research (NIHR). The views expressed are those of the author and not necessarily those of the NHS, the NIHR or the Department of Health.